

Unremovable ‘removable’ singularities

Alexander GOFEN

Dedicated to the memory of Prof. Michael Lidov

ABSTRACT. The article attempts to answer the question why the so called ‘removable’ or ‘regular’ singularities in certain analytic functions cannot be removed. This problem may be understood in the frame of the generalized *elementary functions* (i.e. functions defined as solutions of explicit rational Ordinary Differential Equations). Along with several known examples, the article produces a family of infinitely many functions having regular singularities. There are formulated also two open questions.

1. Introduction

The concept of *removable* (or *regular*) singularities emerges when an analytic function $x(t)$ is presented either as a formula, or as a solution of an Initial Value Problem (IVP) for ODEs, invalid at an isolated point, say $t = 0$, yet valid in its neighborhood. If by convention the proper value is assigned to $x(t)$ at $t = 0$, the function at this point becomes holomorphic, so that its ‘seeming singularity’ is ‘removed’. That is, the singularity contained in the formula or the equations defining the function, does not necessarily belong to this function: for example

$$x(t) = \frac{+\sqrt{1+t}-1}{t}, \quad x|_{t=0} = \frac{1}{2}.$$

As a solution of the polynomial equation $tx^2+2x-1=0$, the function $x(t)$ does not have another algebraic or rational non-singular representation at $t=0$. However $x(t)$ satisfies a regular ODE at $t=0$

$$x' = -\frac{x^2}{2tx+2}, \quad x|_{t=0} = \frac{1}{2}$$

(and some singular ODEs too).

An entire function $x = te^t$ is represented via a regular formula. Still, ODEs defining it may be either singular or regular at $t=0$ (Item 10, Table 1).

However there exist functions for which all currently known formulas or ODEs have a singularity at an isolated point, even though the functions themselves are

1991 *Mathematics Subject Classification*. 34A09, 34A34, 34A12 .

Key words and phrases. ODE, Removable singularities, Regular singularities.

This paper owes a lot to (hot) discussions with Harley Flanders. The beautiful example $x(t) = \cos \sqrt{t}$ is by courtesy of George Bergman.

holomorphic at this point. The examples of such functions are $x(t) = \frac{e^t-1}{t}$, and $x(t) = \cos \sqrt{t}$, and the solution of the IVP

$$tx'' - x = 0; \quad x|_{t=0} = 0; \quad x'|_{t=0} = 1,$$

and each of functions 1-7, Table 1.

Is it possible that regular ODEs representing these functions exist, but are not yet known? This question makes sense only if we specify in which class of equations we are looking for the answer. If the right hand sides of the ODEs are allowed to be any analytic functions, the answer is trivial: Just introduce notation for a new entire analytic function and its derivative.

We consider a subclass of analytic functions called generalized *elementary* functions (first introduced by R. Moore [1]). This class widens the conventionally defined (by Liouville) elementary functions to include practically all functions used in applications. In simplest terms, generalized *elementary* functions [2] are those which may be defined as solutions of IVPs for explicit ODEs having rational right hand sides regular at the initial point. The goal is to prove that $x(t) = \frac{e^t-1}{t}$ and several other functions (Items 1-7, Table 1) cannot satisfy any rational regular ODE at $t = 0$.

All throughout this paper functions and solutions of ODEs are considered as analytic functions in complex space \mathbf{C} .

2. Polynomial ODEs having the same solution

If we are given a polynomial ODE having a solution $x(t)$, it is possible to obtain a non-trivial family of polynomial ODEs (not necessarily just multiplied by a non-zero factor), still having the same solution $x(t)$. We are particularly interested in the case when all derivatives of $x(t)$ are rational at $t = 0$.

LEMMA 1. *Let an analytic function $x(t)$ at the neighborhood of $t = 0$ satisfy a nontrivial polynomial ODE*

$$(2.1) \quad F(t, x, x', \dots, x^{(m)}) = \sum_{k=1}^q a_k t^{\alpha_k} x^{\beta_k} (x')^{\gamma_k} \dots (x^{(m)})^{\omega_k} = 0,$$

$$(2.2) \quad x^{(m)} \Big|_{t=0} = r_m, \quad m = 0, 1, 2, \dots$$

with complex coefficients a_k (k is omitted at power indexes α_k, β_k, \dots). Then this $x(t)$ also satisfies infinitely many polynomial ODEs, such that their coefficients are solutions of a special linear algebraic system 2.4.

In particular, if all derivatives $x^{(m)} \Big|_{t=0}$ are rational, there exists a polynomial ODE with rational coefficients satisfied by $x(t)$.

of the system. This matrix (and the linear system) is infinite only in the number of rows (equations). Only finite number of them are linearly independent. Let the maximal number of linearly independent equations 2.4 be $p > 0$, $p < q$. Therefore there must exist p independent variables with a nonzero sub-determinant corresponding to them, and $q - p$ dependent variables. Among b_1, b_2, \dots, b_q , consider p those which are independent, and assign them rational values. Then the remaining dependent variables must all be rational too (as ratios of sub-determinants of matrix M , whose all elements are rational numbers). The obtained rational coefficients b_1, b_2, \dots, b_q generate the polynomial G_0 having the solution $x(t)$, which completes the proof. \square

EXAMPLE 1. *As an illustration, consider an analytic element $x^{(m)}|_{t=0} = m!$, $m = 0, 1, 2, \dots$ (representing $x = \frac{1}{1-t}$ indeed), and an implicit polynomial equation*

$$(2.6) \quad Ax^2 + Bxt + Cx't + Dx + Ex' + F = 0$$

whose coefficients A, B, \dots are to be determined. By differentiation and substitution of the initial values obtain

$$\begin{aligned} A & & + D & + E & + F = 0 \\ A(m+1)! + Bm! + Cmm! + Dm! + E(m+1)! & = 0, \quad m = 1, 2, \dots \end{aligned}$$

The general solution of this system

$$B = -A - D - E, \quad C = -A - E, \quad F = -A - D - E$$

delivers infinitely many solutions. In particular, the three solutions below exemplify different polynomial equations all satisfied by $x(t)$:

$E = 0, A = 0, D = 1$	$E = 1, A = -1, D = 0$	$E = 1, A = 0, D = -1$
$B = -1, C = 0, F = -1$	$B = 0, C = 0, F = -1$	$B = 0, C = -1, F = 0$
$x - xt - 1 = 0$	$x' - x^2 = 0$	$x' - x't - x = 0$

3. No regular representation for $\frac{e^t-1}{t}$

We deal with the entire function

$$x(t) = \frac{e^t - 1}{t}, \quad x|_{t=0} = 1.$$

It is easily checked that

$$(3.1) \quad x^{(m)}|_{t=0} = \frac{1}{m+1}, \quad m = 0, 1, 2, \dots$$

THEOREM 1. *The function $x(t)$ cannot be a solution of any non-trivial, implicit, polynomial ODE*

$$F(t, x, x', \dots, x^{(m)}) = 0$$

with integer coefficients in the corresponding polynomial

$$F(T, X_0, X_1, \dots, X_m),$$

having

$$\frac{\partial F}{\partial X_m} \Big|_{t=0} \neq 0.$$

PROOF. Denote

$$(3.2) \quad F_0(t, x, \dots, x^{(m)}) = F = \sum a_k t^\alpha x^\beta (x')^\gamma \dots (x^{(m)})^\omega = 0$$

where a_k are integers (k is omitted at power indexes α_k, β_k, \dots).

Repeatedly differentiate relation 3.2, denoting the result of N differentiations by

$$F_N(t, x, \dots, x^{(m)}, \dots, x^{(m+N)}) = \frac{d^N}{dt^N} F_0(t, x, \dots, x^{(m)}).$$

Prove by the induction, that in each of polynomials F_N the highest derivative $x^{(m+N)}$ appears only in one expression always with the same factor $\frac{\partial F_0}{\partial X_m}$. Observe, that

$$\begin{aligned} F_1 &= \frac{d}{dt} F_0(t, x, \dots, x^{(m)}) = \frac{\partial F_0}{\partial X_m} x^{(m+1)} + Q_0(t, x, \dots, x^{(m)}); \\ F_2 &= \frac{d}{dt} F_1(t, x, \dots, x^{(m+1)}) = \frac{\partial F_1}{\partial X_{m+1}} x^{(m+2)} + Q_1(t, x, \dots, x^{(m+1)}) = \\ &= \frac{\partial F_0}{\partial X_m} x^{(m+2)} + Q_1(t, x, \dots, x^{(m+1)}). \end{aligned}$$

Assuming

$$\begin{aligned} F_N &= \frac{\partial F_0(t, x, \dots, x^{(m)})}{\partial X_m} x^{(m+N)} + Q_{N-1}(t, x, \dots, x^{(m+N-1)}); \\ &\quad \frac{\partial F_{N-1}(t, x, \dots, x^{(m+N-1)})}{\partial X_{m+N-1}} = \frac{\partial F_0}{\partial X_m} \end{aligned}$$

to be true for N , obtain

$$\begin{aligned} F_{N+1} &= \frac{d}{dt} F_N(t, x, \dots, x^{(m+N)}) = \frac{\partial F_N}{\partial X_{m+N}} x^{(m+N+1)} + Q_N(t, x, \dots, x^{(m+N)}) = \\ &= \frac{\partial}{\partial X_{m+N}} \left(\frac{\partial F_0}{\partial X_m} \underbrace{x^{(m+N)}}_{\text{the only occurrence of } x^{(m+N)} \text{ in } F_N} + Q_{N-1}(t, x, \dots, x^{(m+N-1)}) \right) x^{(m+N+1)} + \\ &\quad + Q_N(t, x, \dots, x^{(m+N)}) = \frac{\partial F_0}{\partial X_m} x^{(m+N+1)} + Q_N(t, x, \dots, x^{(m+N)}). \end{aligned}$$

Observe that the polynomials F_N have integer coefficients. By the condition of this Theorem, $\frac{\partial F_0}{\partial X_m} \Big|_{t=0} = A \neq 0$. As $x^{(k)}|_{t=0} = \frac{1}{k+1}$, the value A is rational.

Multiply F_0 by a proper integer to clear all denominators so that value $\frac{\partial F_0}{\partial X_m} \Big|_{t=0} = A$ becomes an integer. Then the equation for F_N takes the form:

$$(3.3) \quad F_N = \frac{A}{m+N+1} + Q_{N-1}(t, x, \dots, x^{(m+N-1)}) = 0$$

With growing N , the denominator $m+N+1$ will become greater than A , and then it will reach some prime $p = m+N+1$ so that $\frac{A}{p}$ is a fraction in the lowest terms. All the remaining terms in $Q_{N-1}(t, x, \dots, x^{(m+N-1)})$ must be integers or fractions, whose denominators contain primes less than p . Thus the isolated fraction $\frac{A}{p}$ and $Q_{N-1}(t, x, \dots, x^{(m+N-1)})$ cannot cancel, which is impossible, proving the Theorem. \square

Unlike the previous, the next theorem deals with ODEs having complex (non-integer) coefficients.

THEOREM 2. *The function $x(t)$ cannot be a solution of any non-trivial, implicit, polynomial ODE with complex coefficients*

$$(3.4) \quad F(t, x, x', \dots, x^{(m)}) = 0$$

having

$$(3.5) \quad \left. \frac{\partial F}{\partial X_m} \right|_{t=0} \neq 0.$$

PROOF. Assume the opposite, that ODE 3.4 has $x(t)$ as a solution in a neighborhood of $t = 0$.

Step 0: From complex to real coefficients. Observe, that $x(t)$ and all its derivatives satisfying polynomial 3.4, are real-valued functions on the real axis. Assume therefore the coefficients of polynomial 3.4 are real.

Step 1: From irrational to rational coefficients. According to Lemma 1, $x(t)$ must satisfy infinitely many nontrivial polynomial ODEs with rational coefficients b_1, b_2, \dots, b_q , obtainable as solutions of the linear algebraic equation 2.4. The coefficients a_1, a_2, \dots, a_q of 3.4 are the solutions of linear system 2.4 too. Among them consider the independent ones a_k , and choose their rational approximation b_k so close to a_k , that for the modified polynomial G_0 (Lemma 1, equation 2.5) corresponding to the complete set of rational coefficients b_1, b_2, \dots, b_q , condition 3.5 still holds. To not complicate notation, assume that the given equation 3.4 already has all rational coefficients.

Step 3: Apply a proper integer factor to the polynomial equation 3.4 (having rational coefficients) to clear all denominators. Now $x(t)$ satisfies a polynomial equation with integer coefficients — impossible, according to Theorem 1, which proves this theorem. \square

COROLLARY 1. *The function $x(t)$ cannot be a solution of an IVP for any explicit rational ODE*

$$(3.6) \quad x^{(m+1)} = \frac{P(t, x, x', \dots, x^{(m)})}{Q(t, x, x', \dots, x^{(m)})}$$

having the denominator

$$(3.7) \quad Q|_{t=0} \neq 0,$$

nor indeed it can be a solution of an IVP for any explicit polynomial ODE

$$x^{(m+1)} = P(t, x, x', \dots, x^{(m)}).$$

The proof of this corollary relies on the following

LEMMA 2. *The implicit polynomial ODE 3.4 non-singular at $t = 0$ (Condition 3.5) and the explicit rational ODE 3.6 with a nonzero denominator (Condition 3.7) converts into each other.*

PROOF. Really, in a rational ODE 3.6 written as a polynomial equation

$$F = x^{(m+1)}Q(t, x, x', \dots, x^{(m)}) - P(t, x, x', \dots, x^{(m)}) = 0$$

derivative $\frac{\partial F}{\partial X_{n+1}} \Big|_{t=0} = Q|_{t=0} \neq 0$. Inversely, if a polynomial ODE 3.4 is given, apply $\frac{d}{dt}$

$$\frac{\partial F}{\partial T} + \frac{\partial F}{\partial X} x' + \dots + \frac{\partial F}{\partial X_{m-1}} x^{(m)} + \frac{\partial F}{\partial X_m} x^{(m+1)} = 0$$

and obtain a rational ODE relying on condition 3.5

$$x^{(m+1)} = - \frac{\frac{\partial F}{\partial T} + \frac{\partial F}{\partial X} x' + \dots + \frac{\partial F}{\partial X_{m-1}} x^{(m)}}{\frac{\partial F}{\partial X_m}}$$

□

PROOF. (The Corollary). Assume that the rational ODE 3.6 exists under condition 3.7. According to the Lemma, rational ODE 3.6 converts to the polynomial one. That is impossible according to Theorem 2, which proves this corollary. □

4. Other functions having no regular representation

The method of proof in Theorem 1 applies not only to $x(t)$ having expansion 3.1, but also to infinitely many other analytic functions defined by a variety of expansions (Examples 2-7, Table 1).

COROLLARY 2. Let $H(n) \neq 0$ be an integer-valued function such that the maximal prime $p \leq n$ occurs among the factors of $H(n)$, and let $G(n)$ be an integer-valued function, whose factors do not exceed n . Then the statement of Theorem 1 holds also for functions defined by an analytic element

$$x^{(n)}|_{t=0} = \begin{cases} \frac{G(n-1)}{H(n)} & \text{for infinitely many prime values of } n \\ 0 & \text{otherwise} \end{cases}$$

PROOF. Reconsider equation 3.3 in Theorem 1, which takes the form

$$(4.1) \quad F_N = \frac{AG(m+N-1)}{H(m+N)} + Q_{N-1}(t, x, \dots, x^{(m+N-1)}) = 0.$$

Choose such a big $n = m + N$, that n is prime, $n > A$. Then $\frac{AG(m+N-1)}{H(m+N)}$ is a fraction in the lowest terms, cancellation in equation 4.1 is impossible, proving this corollary. □

It is easy to see that Examples 2-7, Table 1, meet the condition of Corollary 2. Two more examples (not in the Table), defined by their expansion at one point only, also do: $x^{(n)}|_{t=0} = \frac{1}{n!}$, and $x^{(n)}|_{t=0} = \frac{1}{n^n}$. (Other representations of these entire analytic functions are not known).

5. Discussion

Another proof of Theorem 2 belongs to H. Flanders [3,4]. Moreover, he proved that among ODEs of *first* order defining $x = \frac{e^t - 1}{t}$, the known ODEs

$$(5.1) \quad \begin{aligned} x' = R(x) &= \frac{tx - x + 1}{t} \\ P(t, x, x') &= tx' - tx + x - 1 = 0 \end{aligned}$$

are unique in the sense, that any implicit first order polynomial ODE divides by P , while any explicit rational first order ODE reduces to R .

Table 1 summarizes the functions considered in the article. Items (1-7) have no regular representation. Formulas for functions (8,9) are regular at $t = 0$: they are entered into the Table for comparison only. (There exist both regular and singular ODEs for function (8)). We do not know any non-singular rational ODE for the Bessel functions (Item 11), nor is Corollary 2 applicable to them.

5.1. Taylor expansions for elementary functions. Although Theorem 2 and Corollaries 1, 2 for functions (1-7) in Table 1 are about certain specialty of the point $t = 0$ in these functions, it is not yet known whether these functions are *non-elementary* at this isolated point. In order to prove it, a stronger theorem should be established (see the *Proposition* in the next section). We can only suspect that $x(t)$ is possibly non-elementary at $t = 0$. If so, then any system of rational ODEs satisfied by $x(t)$ must be singular, so that specialty of the point $t = 0$ in $x(t)$ is ‘unremovable’ in the class of elementary functions.

	Functions	ODEs	Derivatives at $t = 0$
1	$x = \frac{e^t - 1}{t}$	$x' = \frac{tx - x + 1}{t}$	$x^{(n)} = \frac{1}{n+1}$
2	$x = \frac{\sin t}{t}$ $y = \cos t$ $z = \sin t$	$x' = \frac{y-x}{t}$ $y' = -z$ $z' = y$	$x^{(n)} = \frac{(-1)^n}{2n+1}$ even n , or 0 $y^{(n)} = (-1)^n$ even n , or 0 $z^{(n)} = (-1)^{n+1}$ odd n , or 0
3	$x = \frac{\cos t - 1}{t^2}$ $y = \cos t$ $z = \sin t$	$x' = \frac{2-2y-tz}{t^3}$ $y' = -z$ $z' = y$	$x^{(n)} = \frac{(-1)^{n+1}}{(2n+1)(2n+2)}$ even n , or 0 $y^{(n)} = (-1)^n$ even n , or 0 $z^{(n)} = (-1)^{n+1}$ odd n , or 0
4	$x = \cos \sqrt{t}$ $y = \sin \sqrt{t}$ $z = \sqrt{t}$	$x' = -\frac{yz}{2t}$ $y' = \frac{xz}{2t}$ $z' = \frac{z}{2t}$	$x^{(n)} = (-1)^n \frac{n!}{(2n)!}$ singular singular
5	$x = \frac{\cos \sqrt{t} - 1}{t}$ $u = \cos \sqrt{t}$ $v = \sin \sqrt{t}$ $z = \sqrt{t}$	$x' = \frac{-vz - 2u + 2}{2t^2}$ $u' = -\frac{vz}{2t}$ $v' = \frac{uz}{2t}$ $z' = \frac{z}{2t}$	$x^{(n)} = (-1)^{n+1} \frac{(n-1)!}{(2n)!}$ $u^{(n)} = (-1)^n \frac{n!}{(2n)!}$ singular singular
6	$tx'' - x = 0$	$x'' = \frac{x}{t}$	$x^{(n)} = \frac{1}{(n-1)!}$, $n \geq 1$, $x(0) = 0$
7	$x = \frac{\ln(t+1)}{t}$	$x' = \frac{1-tx-x}{t(t+1)}$	$x^{(n)} = \frac{(-1)^{n+1} n!}{n+1}$
8	$x = \ln(t+1)$	$x' = \frac{1}{t+1}$	$x^{(n)} = (-1)^{n-1} (n-1)!$
9	$x = e^t$	$x' = x$	$x^{(n)} = 1$
10	$x = te^t$	$x' = \frac{x}{t} + x$ $x'' = 2x' - x$	$x^{(n)} = n$
11	Bessel functions J_p , $p = 0, 1, 2, \dots$	$x'' = \frac{-tx' - (t^2 - p^2)x}{t^2}$	$x^{(n)} = \frac{(-1)^k C_{2k+p}^k}{2^{2k+p}}$, $n = 2k + p$ or 0

Table 1. Summary of functions, ODEs defining them, and their n -order derivatives

Elementary functions represent practically all functions used in applications, and they are elementary (almost) at all points of their holomorphy. Yet their Taylor expansions have certain specialty, distinguishing them from non-elementary functions: their Taylor coefficients are obtainable via a fixed number of *explicit* formulas of Automatic Differentiation (AD), corresponding to a system of *explicit* rational ODEs and algebraic relations [2]. Systems of *implicit* rational ODEs and

implicit algebraic relations are considered by Nedialkov and Pryce [5]. Generally, an expansion generated by an arbitrary recursive formula or algorithm may not be expected to represent a function being elementary at this or other points.

5.2. Open statements. The method of proof of Theorem 2 for an n -order ODE is not applicable to *systems* of ODEs, leaving open the following

PROPOSITION 1. *An entire function*

$$x(t) = \frac{e^t - 1}{t}, \quad x^{(m)}|_{t=0} = \frac{1}{m + 1}, \quad m = 0, 1, 2, \dots$$

at the point $t = 0$ cannot be a solution of an IVP for any system of rational ODEs

$$(5.2) \quad \begin{aligned} x' &= \frac{P_1(t, x, y, z, \dots)}{Q_1(t, x, y, z, \dots)} \\ y' &= \frac{P_2(t, x, y, z, \dots)}{Q_2(t, x, y, z, \dots)} \\ &\dots \end{aligned}$$

whose all denominators $Q_i|_{t=0} \neq 0$, nor indeed it can be a solution of an IVP for any system of explicit polynomial ODEs

$$\begin{aligned} x' &= P_1(t, x, y, z, \dots) \\ y' &= P_2(t, x, y, z, \dots) \\ &\dots \end{aligned}$$

If proved, this Proposition would establish existence of a new type of special points in elementary analytic functions (along with Poles, Branching, and Essential singularities).

Another open statement (which, if proved, would solve Proposition 1), is the following

CONJECTURE 1. *Consider an IVP for a system of rational ODEs 5.2 with nonzero denominators at $t = 0$, so that it has a unique regular solution in a neighborhood of $t = 0$. In particular, derivatives $x^{(n)}|_{t=0} = a_n, n = 0, 1, 2, \dots$. Then there exists an explicit rational ODE of order $n \geq 1$*

$$x^{(n+1)} = \frac{F(t, x, \dots, x^{(n)})}{G(t, x, \dots, x^{(n)})}; \quad x^{(k)}|_{t=0} = a_k, \quad k = 1, 2, \dots, n$$

whose denominator $G|_{t=0} \neq 0$, having $x(t)$ as a unique solution.

The Conjecture claims convertibility of an explicit first order system of rational ODEs into one explicit rational ODE of order n . (The opposite conversion is well known and trivial).

6. References

[1] Moore, R. E., 1966, Interval analysis, pp. 107-130, Prentice-Hall, Englewood Cliffs, N.J.
 [2] Gofen, A., 2002, ODEs and redefining the concept of elementary functions, In: P.M.A. Sloot et al. (Eds), Computational science - ICCS 2002, LNCS 2329, pp. 1000-1009, Amsterdam, Springer
 [3] Flanders, H., 2006, Solutions of ODEs with removable singularities, In: M. Bucker, G. Corliss et al. (Eds), Automatic Differentiation: Applications, Theory,

and Implementations. Lecture notes in Computational science and engineering, pp. 35 - 45, Springer-Verlag, Berlin.

[4] Flanders, H., 2007, Functions not satisfying implicit polynomial ODE, J. Differential Equations, vol. 240, issue 1, September, pp. 164-171.

[5] Nedialkov N.S., J.D. Pryce, 2005, Solving differential-algebraic equations by Taylor series, BIT 45, No.3, 561-591.

THE SMITH-KETTLEWELL EYE RESEARCH INSTITUTE, 2318 FILLMORE ST., SAN FRANCISCO, CA 94115, USA

E-mail address: galex@ski.org

URL: <http://TaylorCenter.org>